

POST-TEST

Consensus or Controversy, Issue 1: Clinical Investigators Provide Their Perspectives on Controversial Issues in the Use of Chemotherapy in the Management of Hormone Sensitive Metastatic Prostate Cancer (mPCa)

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

1. The systemic treatment approach preferred by the majority of the faculty for a 62-year-old man with a good performance status and extensive bone and liver metastases but only minimal complaints of pain is _____.

 - a. Androgen deprivation therapy (ADT) alone
 - b. ADT and docetaxel
 - c. Docetaxel alone
2. For a 62-year-old man with a good performance status and extensive bony metastases but only minimal complaints of pain, all of the faculty would administer ADT with docetaxel-based therapy.

 - a. True
 - b. False
3. For an 80-year-old man with extensive bony metastases and recent complaints of pain, fatigue and weight loss (PS = 2) that are believed to be cancer related, the majority of the faculty would recommend ADT with docetaxel.

 - a. True
 - b. False
4. Regardless of patient age, the majority of investigators would administer _____ to an asymptomatic man with a good performance status and 3 isolated rib metastases.

 - a. ADT alone
 - b. ADT and docetaxel
 - c. Docetaxel alone