



POST-ASH Issue 5, 2015

**Updated Results from the Phase
III APL0406 Trial of ATRA/
Arsenic Trioxide versus ATRA/
Chemotherapy in Nonhigh-Risk APL**

For more visit ResearchToPractice.com/5MJCASH2015

Research
To Practice®

CME INFORMATION

OVERVIEW OF ACTIVITY

Each year, thousands of clinicians, basic scientists and other industry professionals sojourn to major international oncology conferences, like the American Society of Hematology (ASH) annual meeting, to hone their skills, network with colleagues and learn about recent advances altering state-of-the-art management in hematologic oncology. As such, these events have become global stages where exciting science, cutting-edge concepts and practice-changing data emerge on a truly grand scale. This massive outpouring of information has enormous benefits for the hematologic oncology community, but the truth is it also creates a major challenge for practicing oncologists and hematologists.

Although original data are consistently being presented and published, the flood of information unveiled during a major academic conference is unprecedented and leaves in its wake an enormous volume of new knowledge that practicing oncologists must try to sift through, evaluate and consider applying. Unfortunately and quite commonly, time constraints and an inability to access these data sets leave many oncologists struggling to ensure that they're aware of crucial practice-altering findings. This creates an almost insurmountable obstacle for clinicians in community practice because they are not only confronted almost overnight with thousands of new presentations and data sets to consider but they are also severely restricted in their ability to review and interrogate the raw findings.

To bridge the gap between research and patient care, this CME activity will deliver a serial review of the most important emerging data sets on up-front and salvage therapeutic options and modalities for the evaluation of treatment response in newly diagnosed and relapsed or refractory acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL), acute promyelocytic leukemia (APL) and myelodysplastic syndromes (MDS) from the latest ASH meeting, including expert perspectives on how these new evidence-based concepts may be applied to routine clinical care. This activity will assist medical oncologists, hematologists, hematology-oncology fellows and other healthcare professionals in the formulation of optimal clinical management strategies and the timely application of new research findings to best-practice patient care.

LEARNING OBJECTIVES

- Evaluate the results of the confirmatory Phase II BLAST study of blinatumomab in patients with relapsed/refractory ALL, and consider this information when developing treatment plans for these patients.
- Analyze the efficacy of a more intensive pediatric chemotherapy regimen for older adolescents and young adults with newly diagnosed B- or T-precursor ALL, and determine the feasibility of this approach for patients with ALL in this age group.
- Compare and contrast the benefits and risks reported in the Phase III APL0406 trial of all-trans retinoic acid (ATRA) with arsenic trioxide versus ATRA with chemotherapy, and consider the potential therapeutic benefit of a chemotherapy-free regimen for patients with newly diagnosed nonhigh-risk APL.
- Determine the clinical benefit seen with the addition of the multikinase inhibitor sorafenib to standard primary induction and consolidation therapy for younger patients with newly diagnosed AML.
- Assess the efficacy and tolerability profile of the novel agent vosaroxin combined with cytarabine from the Phase III VALOR trial in patients with relapsed or refractory AML.
- Examine the impact of lenalidomide therapy on the achievement of transfusion independence in red blood cell transfusion-dependent patients with lower-risk MDS without del(5q).

ACCREDITATION STATEMENT

Research To Practice is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

CREDIT DESIGNATION STATEMENT

Research To Practice designates this enduring material for a maximum of 1.5 *AMA PRA Category 1 Credits™*. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

HOW TO USE THIS CME ACTIVITY

This CME activity contains slides and edited commentary. To receive credit, the participant should review the slide presentations, read the commentary, complete the Post-test with a score of 75% or better and fill out the Educational Assessment and Credit Form located at ResearchToPractice.com/5MJCASH2015/5/CME.

CONTENT VALIDATION AND DISCLOSURES

Research To Practice (RTP) is committed to providing its participants with high-quality, unbiased and state-of-the-art education. We assess potential conflicts of interest with faculty, planners and managers of CME activities. Real or apparent conflicts of interest are identified and resolved through a conflict of interest resolution process. In addition, all activity content is reviewed by both a member of the RTP scientific staff and an external, independent physician reviewer for fair balance, scientific objectivity of studies referenced and patient care recommendations.

FACULTY — The following faculty (and their spouses/partners) reported real or apparent conflicts of interest, which have been resolved through a conflict of interest resolution process:

Mikkael A Sekeres, MD, MS
Professor of Medicine
Director, Leukemia Program
Cleveland Clinic Taussig Cancer Institute
Cleveland, Ohio

Advisory Committee: Amgen Inc, Boehringer Ingelheim Pharmaceuticals Inc, Celgene Corporation.

EDITOR — Dr Love is president and CEO of Research To Practice, which receives funds in the form of educational grants to develop CME activities from the following commercial interests: AbbVie Inc, Amgen Inc, Astellas Scientific and Medical Affairs Inc, AstraZeneca Pharmaceuticals LP, Aveo Pharmaceuticals, Bayer HealthCare Pharmaceuticals, Biodesix Inc, Boehringer Ingelheim

Pharmaceuticals Inc, Boston Biomedical Pharma Inc, Bristol-Myers Squibb Company, Celgene Corporation, Clovis Oncology, Daiichi Sankyo Inc, Dendreon Corporation, Eisai Inc, Exelixis Inc, Foundation Medicine, Genentech BioOncology, Genomic Health Inc, Gilead Sciences Inc, Incyte Corporation, Janssen Biotech Inc, Jazz Pharmaceuticals Inc, Lilly, Medivation Inc, Merck, Myriad Genetic Laboratories Inc, NanoString Technologies, Novartis Pharmaceuticals Corporation, Novocure, Onyx Pharmaceuticals, an Amgen subsidiary, Pharmacyclics Inc, Prometheus Laboratories Inc, Regeneron Pharmaceuticals, Sanofi, Seattle Genetics, Sigma-Tau Pharmaceuticals Inc, Sirtex Medical Ltd, Spectrum Pharmaceuticals Inc, Taiho Oncology Inc, Takeda Oncology, Teva Oncology, Tokai Pharmaceuticals Inc and VisionGate Inc.

RESEARCH TO PRACTICE STAFF AND EXTERNAL REVIEWERS —

The scientific staff and reviewers for Research To Practice have no real or apparent conflicts of interest to disclose.

This educational activity contains discussion of published and/or investigational uses of agents that are not indicated by the Food and Drug Administration. Research To Practice does not recommend the use of any agent outside of the labeled

indications. Please refer to the official prescribing information for each product for discussion of approved indications, contraindications and warnings. The opinions expressed are those of the presenters and are not to be construed as those of the publisher or grantors.

This activity is supported by educational grants from Bristol-Myers Squibb Company, Celgene Corporation, Incyte Corporation, Onyx Pharmaceuticals, an Amgen subsidiary, Seattle Genetics and Takeda Oncology.

Hardware/Software Requirements:

A high-speed Internet connection

A monitor set to 1280 x 1024 pixels or more

Internet Explorer 7 or later, Firefox 3.0 or later, Chrome,

Safari 3.0 or later

Adobe Flash Player 10.2 plug-in or later

Adobe Acrobat Reader

(Optional) Sound card and speakers for audio

Last review date: May 2015

Expiration date: May 2016

To go directly to slides and commentary for this issue, [click here](#).

Not long after this year's American Society of Hematology (ASH) meeting, we gathered 6 clinical investigators for our first ever think tank focused exclusively on leukemias and myelodysplastic syndromes (MDS). Although a lot of the excitement during this closed recording session centered on new agents and therapies — particularly the explosion of encouraging clinical research in acute lymphoblastic leukemia (ALL) with both CAR T-cell immunotherapy and the bispecific T-cell engager antibody blinatumomab (see below) — it was also fascinating to hear that an older drug that has sometimes gotten a bad rap — sorafenib (sor) — may have a new role as part of up-front treatment for acute myeloid leukemia (AML).

ALL is an uncommon disease that many oncologists appropriately triage to tertiary centers, but AML — particularly in elderly patients — is an important part of general oncology practice. As such, even though the randomized, Phase II trial of sor presented at ASH as a plenary was not the “home run” that we are beginning to see more frequently with immunotherapy in many diseases, from a practical clinical perspective the study findings may be among the most important in any cancer this year.

For that reason, we lead off this year's acute leukemia/MDS ASH summary by focusing on that work. But as always, we also created teaching slide sets and obtained perspectives from a noted clinical investigator — in this case Mikkael Sekeres — for a number of the most important presentations, which are outlined below:


AML

Sor up front in AML

About 20% of patients with AML have activating mutations in the FMS-like tyrosine kinase 3 (FLT3), and a number of FLT3 inhibitors are in various states of development. Sor targets this kinase, among others, and for that reason this [Phase II placebo-controlled German trial](#) evaluated the addition of sor to standard induction and consolidation treatment (followed by maintenance with sor) in 267 adult patients with

Leukemias and Myelodysplastic Syndromes Think Tank Faculty January 29, 2015

Jennifer R Brown, MD, PhD
Hagop M Kantarjian, MD
Charles A Schiffer, MD
B Douglas Smith, MD
David P Steensma, MD
Wendy Stock, MD, MA



Mikkael A Sekeres, MD, MS

newly diagnosed AML age 60 years and younger. Importantly, individuals both with and without FLT3-internal tandem duplication (ITD) mutations were eligible for and enrolled in the study. With a median follow-up of 3 years, the trial met its primary endpoint of event-free survival (EFS), demonstrating a significant improvement in favor of sor (median EFS 20.5 months versus 9.2 months, $p = 0.013$). Interestingly and quite unexpectedly, there was a suggestion that the benefit was, if anything, more impressive in patients without FLT3-ITD mutations. The one sticking point is, to date there is no overall survival (OS) advantage, which is concerning to Dr Sekeres and has informed his current decision not to use up-front sor outside of a clinical trial.

However, at the think tank the reaction to these data was quite different, as Dr Hagop Kantarjian noted that since 2005, he and his MD Anderson colleagues have been routinely using FLT3 inhibitors in patients with FLT3-ITD mutations and that the outcomes appear indirectly to be improved compared to earlier series. The think tank faculty speculated on possible biologic explanations for these compelling findings, including the presence of other

kinase targets or inhibition of wild-type FLT3 kinase activation, but most seemed to agree that these new data at the very least deserve careful consideration in patients with and without these abnormalities. To further drive home this point, Dr Kantarjian made an impassioned plea for “leukemia doctors to act more like those focused on solid tumors” and seek small research advances that, when coupled together, create a major positive effect for patients, as seen, for example, in renal cell carcinoma.


VALOR trial of vosaroxin

VALOR is a large, **international Phase III study** evaluating cytarabine with or without vosaroxin, a first-in-class anticancer quinolone derivative, in patients with relapsed/refractory AML. On the surface things look straightforward, as the trial did not reach its primary endpoint of improved OS. However, the data also demonstrated that complete remission rates were improved with vosaroxin/cytarabine, and a preplanned survival subgroup analysis censoring patients at allogeneic transplant showed a statistically significant 1.4-month advantage (hazard ratio 0.83, $p = 0.02$). Dr Sekeres is not convinced these improvements are clinically meaningful, but Dr Kantarjian — whose group has done a lot of this research — believes vosaroxin has important value and should be made available to clinicians.

MDS

Lenalidomide (len) in non-del(5q) disease

While the role of this immunomodulatory agent is well established and approved in patients with del(5q), mainly for management of anemia, a prior Phase II trial



[Click here to see Dr Kantarjian's comment](#)

suggested clear-cut benefit in non-del(5q) disease. [This Italian Phase III study](#) almost duplicated the results seen in the Phase II effort and demonstrated a ≥ 56 -day transfusion independence rate of 27% with len compared to a 2.5% rate with placebo. These findings will undoubtedly lead clinicians to want to use this drug in this situation, and think tank participant Dr David Steensma endorses this approach. However, he cautions that “platelets need to be at a reasonable level” to use len.

Azacitidine (aza) alone or with len or vorinostat (vor) in higher-risk MDS and chronic myelomonocytic leukemia

At ASH Dr Sekeres presented the first results from the largest prospective study in higher-risk MDS ever conducted in North America — [SWOG-S1117](#) — which demonstrated a modest signal for improvement in disease-related outcomes with the 2 combinations. Unfortunately, a greater likelihood to discontinue treatment due to toxicity was also seen (9% aza, 23% aza/len, 24% aza/vor). In discussing this work, Dr Sekeres pointed out that these data will continue to mature, and he believes it is possible that with better management of side effects, these and other combinations may be successfully incorporated into treatment.

ATRA/arsenic trioxide (AAT) in acute promyelocytic leukemia (APL)

At the 2012 ASH meeting, the initial findings from the landmark Phase III Italian-German APL0406 trial in low/intermediate-risk APL comparing AAT as induction and consolidation to ATRA/idarubicin as induction, consolidation and maintenance therapy grabbed headlines and led many oncologists to change their practices. Dr Sekeres and his Cleveland Clinic group, however, wanted to see more follow-up before following suit. That information came [at this year's meeting](#) as excellent outcomes (now with 254 patients evaluable for response at 3 years) were observed with both therapies, but there appeared to be a suggestion of greater benefit with AAT (complete response [CR] 100% versus 97%; 2-year EFS rate 98% versus 84.9%, $p = 0.0002$; 2-year OS rate 99.1% versus 94.4%, $p = 0.01$). This has now given Dr Sekeres and his group enough supporting evidence to offer the chemotherapy-free AAT combination as standard induction and postremission therapy to patients with low/intermediate-risk APL.

ALL

Treatment for older adolescents and young adults (AYAs)

For years, a fundamental issue in this disease has been whether more intensive pediatric regimens should be used in AYAs. At ASH we saw relevant findings from the single-arm [US Intergroup trial C10403](#) of 296 patients age 16 to 39 who received a pediatric regimen administered by adult hematologist-oncologists. The 2-year EFS of 66% seen in this study represents a significant improvement compared to 34% EFS observed in historical controls, and globally the outcomes, including toxicities, were similar to what has been documented in other prospective international studies of pediatric regimens in AYAs.

As a result of these important findings, Dr Sekeres and the think tank faculty, including Dr Wendy Stock, who presented these data at ASH, all support the use of this approach moving forward both in clinical practice and in trials attempting to integrate new agents. It should also be noted that Dr Kantarjian believes that hyper-CVAD is an equivalent alternative.

CD19-targeted 19-28z CAR-modified autologous T cells in adult patients with relapsed, refractory B-cell ALL

A number of our CME programs have helped chronicle the amazing story of CAR T-cell therapy, and we would be remiss to not provide an update coming out of the year's biggest meeting. As previously mentioned, ALL is the locus where this therapy has taken off, and in San Francisco we saw extended follow-up from a [**Phase I Memorial Sloan Kettering study**](#) in this disease. Of the 22 patients evaluable for response, many of whom had heavily pretreated disease, an impressive 91% (20 patients) achieved CR after CAR T-cell infusion, with 90% (18 patients) of those being MRD-negative. Ten of the 13 transplant-eligible patients subsequently went on to successfully receive an allogeneic hematopoietic cell transplant.

In terms of complications, patients with MRD at the time of treatment did not experience cytokine release syndrome (CRS), and for those with morphologic disease at the time of T-cell infusion, a temporal relationship between serum IL-6 levels and CRS suggests that early intervention with IL-6-directed therapy might be effective in ameliorating related neurologic toxicities.

In commenting on this study, Dr Sekeres cautions that currently CAR T-cell therapy requires specialized administration logistics and the capability to manage potentially challenging cytokine-mediated toxicities. He also questions the long-term durability of response and envisions a future for this approach as a bridge to transplant but is uncertain as to whether CAR T-cell therapy will one day have a role as a stand-alone treatment or as part of induction.

Blinatumomab

At ASH we saw the presentation of the [**Phase II BLAST trial**](#) of 116 patients who were MRD-positive ($\geq 10^{-3}$) after having received at least 3 prior intensive chemotherapy regimens. The MRD CR after 1 cycle of blinatumomab was 78% and did not differ across multiple patient demographics, including those with higher MRD burden. However, the adverse event (AE) profile (mainly related to cytokine release) is not insignificant. Importantly, in this trial serious AEs occurred in 60% of patients, with 2 fatalities.

In discussing the recent FDA accelerated approval of the drug at the think tank, the faculty noted its impressive effectiveness as a salvage therapy but also related the challenges they have faced in managing toxicities. In this regard, Dr Steensma emphasized the role of corticosteroids in mitigating side effects such as fever and impaired mental function. Not surprisingly, a number of current trials combine

blinatumomab with chemotherapy in both the salvage and front-line settings, including a Phase III trial in newly diagnosed ALL ([NCT02003222](#)).

Be on the lookout for the entire think tank program this summer, but next on this series, we talk about new agents in multiple myeloma, particularly the search for the “rituximab of myeloma” that includes a new wave of monoclonal antibodies such as elotuzumab and daratumumab.

Neil Love, MD

[Research To Practice](#)

Miami, Florida

Research To Practice is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Research To Practice designates this enduring material for a maximum of 1.5 *AMA PRA Category 1 Credits™*. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

This activity is supported by educational grants from Bristol-Myers Squibb Company, Celgene Corporation, Incyte Corporation, Onyx Pharmaceuticals, an Amgen subsidiary, Seattle Genetics and Takeda Oncology.

Updated Results from the Phase III APL0406 Trial of ATRA/Arsenic Trioxide versus ATRA/Chemotherapy in Nonhigh-Risk APL

Presentation discussed in this issue

Platzbecker U et al. **Improved outcome with ATRA-arsenic trioxide compared to ATRA-chemotherapy in non-high risk acute promyelocytic leukemia — Updated results of the Italian-German APL0406 trial on the extended final series.** *Proc ASH 2014*; **Abstract 12.**

Slides from a presentation at ASH 2014 and transcribed comments from a recent interview with Mikkael A Sekeres, MD, MS (1/20/15)

Improved Outcome with ATRA-Arsenic Trioxide Compared to ATRA-Chemotherapy in Non-High Risk Acute Promyelocytic Leukemia — Updated Results of the Italian-German APL0406 Trial on the Extended Final Series

Platzbecker U et al.

Proc ASH 2014; Abstract 12.

Research
To Practice®


Background

- Results from the Phase III APL0406 trial showed that the combination of all-trans retinoic acid (ATRA) and arsenic trioxide (ATO) is at least not inferior and possibly superior to standard ATRA and chemotherapy (CHT) in the front-line management of low-to intermediate-risk acute promyelocytic leukemia (APL) (*NEJM* 2013;369(2):111-21):
 - 2-year event-free survival (EFS): 97% (ATRA-ATO) vs 86% (ATRA-CHT); $p < 0.001$ for noninferiority and $p = 0.02$ for superiority
 - 2-year overall survival (OS): 99% (ATRA-ATO) vs 91% (ATRA-CHT); $p = 0.02$
- **Study objective:** To report the updated efficacy results from the extended and final series of 276 patients with newly diagnosed nonhigh-risk APL.

Platzbecker U et al. *Proc ASH* 2014;Abstract 12.

Research
To Practice®

Phase III APL0406 Trial Design



- ATRA: 45 mg/m² per day; ATO: 0.15 mg/kg per day
- Patients on the ATRA-ATO arm received ATRA-ATO induction until complete response (CR), then ATO 5 days/week, 4 weeks on, 4 weeks off for a total of 4 courses and ATRA 2 weeks on, 2 weeks off for a total of 7 courses.
- Patients on the ATRA-CHT arm received the standard ATRA + idarubicin induction followed by 3 cycles of anthracycline-based consolidation together with ATRA and low-dose CHT and ATRA for maintenance.
- **Primary endpoint:** 2-year EFS

Platzbecker U et al. *Proc ASH* 2014;Abstract 12;
LoCoco F et al. *N Engl J Med* 2013;369(2):111-21.

Research
To Practice®

Survival Outcomes

Outcome	ATRA-ATO (n = 122)	ATRA-CHT (n = 132)	p-value
Two-year EFS	98%	84.9%	0.0002
Two-year cumulative incidence of relapse	1.1%	9.4%	0.005
Two-year OS	99.1%	94.4%	0.01

- Median follow-up period = 36 months.
- Four patients died during induction therapy on the ATRA-CHT arm.

Platzbecker U et al. *Proc ASH* 2014;Abstract 12 (Abstract only).

Research
To Practice®

Response to Induction Therapy

	ATRA-ATO (n = 122)*	ATRA-CHT (n = 132)*	p-value
CR rate	100%	97%	0.12

* Number of patients evaluable for response to induction

Platzbecker U et al. *Proc ASH* 2014;Abstract 12 (Abstract only).

Research
To Practice®

Author Conclusions

- The data on this extended cohort demonstrate a significantly augmented survival benefit coupled to a higher antileukemic efficacy provided by ATRA-ATO compared to ATRA-CHT for patients with low- to intermediate-risk APL.
- These results further support ATRA-ATO as the new standard therapy in this clinical setting.

Platzbecker U et al. *Proc ASH 2014*;Abstract 12 (Abstract only).

Research
To Practice®

Investigator Commentary: Updated Results of the APL0406 Trial of ATRA-ATO versus ATRA-CHT in Newly Diagnosed APL

The original results of the APL0406 trial were presented at a plenary session at ASH 2012 and were published in 2013. I believe that the updated results are practice changing. Patients had to have low- to intermediate-risk APL to be eligible. Neither of the treatment arms included cytarabine, which continues to be somewhat controversial in the treatment of APL. The primary endpoint was EFS at 2 years.

When the results were initially presented the median follow-up was 2 years, and we thought the results were provocative but we were eager to see the data after a longer follow-up period. With 3 years of follow-up the CR rate is 100% with ATRA-ATO versus 97% with ATRA-CHT. These results were not statistically different. Four patients died on the ATRA-CHT arm. The CR rate for APL is higher than 90%. Those patients who do not achieve a CR die: There is no middle ground in APL. The 2-year EFS and OS rates were 98% and 99% with ATRA-ATO versus 84.9% and 94.4% with ATRA-CHT. Based on these data, our standard for patients with lower-risk APL at the Cleveland Clinic is induction ATRA-ATO in postremission therapy, and we exclude chemotherapy altogether. We did not make this change after the initial presentation because we were not convinced by the results after just 2 years of follow-up.

Interview with Mikkael A Sekeres, MD, MS, January 20, 2015