POST-TEST

Cases from the Community: Urologic Oncology Investigators Provide Perspectives on the Optimal Management of Prostate Cancer (Webinar Video Proceedings)

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

- 1. Ongoing Phase III trials are evaluating talazoparib in combination with which of the following agents for prostate cancer?
 - a. Enzalutamide
 - b. ¹⁷⁷Lu-PSMA-617
 - c. Darolutamide
 - d. All of the above
- 2. The addition of which of the following to docetaxel and ADT for metastatic hormone-sensitive prostate cancer resulted in a statistically significant improvement in overall survival in the Phase III ARASENS study?
 - a. Apalutamide
 - b. Enzalutamide
 - c. Darolutamide
 - d. Relugolix
- 3. The results of the Phase III PROpel trial of first-line olaparib in combination with abiraterone versus abiraterone alone reported an improvement in radiographic progression-free survival for patients with metastatic castration-resistant prostate cancer (mCRPC) and what homologous recombination repair (HRR) gene mutation status?
 - a. HRR gene mutations
 - b. No HRR gene mutations
 - c. All patients regardless of HRR gene mutation status
 - d. No patients regardless of HRR gene mutation status

- 4. Which of the following any-grade adverse events was most commonly observed in patients with mCRPC receiving ¹⁷⁷Lu-PSMA-617 in the VISION study?
 - a. Bruising
 - b. Thrombocytopenia
 - c. Leukopenia
 - d. Dry mouth
- 5. Which agent resulted in a lower cumulative incidence of major cardiovascular adverse events for patients with advanced prostate cancer in the HERO trial?
 - a. Relugolix
 - b. Leuprolide
 - c. Degarelix
 - d. Sipuleucel-T