## POST-TEST

Oncology Today with Dr Neil Love: Optimizing the Management of Metastatic BRCA-Negative, Triple-Negative Breast Cancer

## THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

- 1. The ongoing Phase III ASCENT-03 study is evaluating sacituzumab govitecan versus treatment of physician's choice in which of the following settings?
  - a. As adjuvant treatment for localized TNBC
  - b. As first-line treatment for mTNBC
  - c. As second-line treatment for mTNBC
  - d. As late-line treatment for mTNBC
- 2. Which of the following Grade 3 adverse events was most commonly reported in patients receiving sacituzumab for previously treated advanced TNBC in the Phase III ASCENT trial?
  - a. Sinus bradvcardia
  - b. Neutropenias
  - c. Peripheral neuropathy
  - d. Nausea
- 3. Recently presented pooled data analysis from pivotal studies of trastuzumab deruxtecan (T-DXd) for patients with HER2-positive metastatic breast cancer (mBC) with brain metastases (BMs) demonstrated which of the following outcomes?
  - a. Significant CNS activity with T-DXd for patients with HER2-positive mBC with treated/stable BMs
  - b. Significant CNS activity with T-DXd for patients with HER2-positive mBC with untreated/active BMs
  - c. Both a and b
  - d. Neither a nor b

- 4. Which of the following receptors is the target of datopotamab deruxtecan (Dato-DXd)?
  - a. NRG1
  - b. MET
  - c. CEACAM5
  - d. TROP2
- 5. Recently presented findings on the combination of Dato-DXd and durvalumab demonstrated robust and durable tumor responses for patients with TNBC in which of the following settings?
  - a. As adjuvant therapy for localized TNBC
  - b. As first-line therapy for unresectable locally advanced/metastatic TNBC
  - c. As second-line therapy for unresectable locally advanced/ metastatic TNBC
  - d. As later-line therapy for unresectable locally advanced/metastatic TNBC