



**POST-ASH** Issue 7, 2014

# **Reinduction of Complete Remissions Using 19-28z CAR T-Cell Therapy in Relapsed B-Cell ALL**

For more visit [ResearchToPractice.com/5MJCASH2014](http://ResearchToPractice.com/5MJCASH2014)

**Research  
To Practice®**

## CME INFORMATION

### OVERVIEW OF ACTIVITY

Each year, thousands of clinicians, basic scientists and other industry professionals sojourn to major international oncology conferences, like the American Society of Hematology (ASH) annual meeting, to hone their skills, network with colleagues and learn about recent advances altering state-of-the-art management in hematologic oncology. As such, these events have become global stages where exciting science, cutting-edge concepts and practice-changing data emerge on a truly grand scale. This massive outpouring of information has enormous benefits for the hematologic oncology community, but the truth is it also creates a major challenge for practicing oncologists and hematologists.

Although original data are consistently being presented and published, the flood of information unveiled during a major academic conference is unprecedented and leaves in its wake an enormous volume of new knowledge that practicing oncologists must try to sift through, evaluate and consider applying. Unfortunately and quite commonly, time constraints and an inability to access these data sets leave many oncologists struggling to ensure that they're aware of crucial practice-altering findings. This creates an almost insurmountable obstacle for clinicians in community practice because they are not only confronted almost overnight with thousands of new presentations and data sets to consider but they are also severely restricted in their ability to review and interrogate the raw findings.

To bridge the gap between research and patient care, this CME activity will deliver a serial review of the most important emerging data sets on chimeric antigen receptor T-cell therapy for leukemia and lymphoma from the latest ASH meeting, including expert perspectives on how these new evidence-based concepts may be applied to routine clinical care. This activity will assist medical oncologists, hematologists, hematology-oncology fellows and other healthcare professionals in the formulation of optimal clinical management strategies and the timely application of new research findings to best-practice patient care.

### LEARNING OBJECTIVES

- Develop an understanding of the mechanism of action of chimeric antigen receptor T-cell therapy, and evaluate the emerging efficacy and safety data with this therapeutic approach under evaluation in the front-line and relapsed/refractory settings for B-cell lymphoma and leukemias.
- Evaluate the benefits and risks of the addition of gemtuzumab ozogamicin to standard chemotherapy and of other emerging agents such as novel FLT3 inhibitors or hypomethylating agents for the treatment of acute myeloid leukemia.

### ACCREDITATION STATEMENT

Research To Practice is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

### CREDIT DESIGNATION STATEMENT

Research To Practice designates this enduring material for a maximum of 2 *AMA PRA Category 1 Credits™*. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

### HOW TO USE THIS CME ACTIVITY

This CME activity contains slides and edited commentary. To receive credit, the participant should review the slide presentations, read the commentary, complete the Post-test with a score of 75% or better and fill out the Educational Assessment and Credit Form located at [ResearchToPractice.com/5MJCASH2014/7/CME](http://ResearchToPractice.com/5MJCASH2014/7/CME).

### CONTENT VALIDATION AND DISCLOSURES

Research To Practice (RTP) is committed to providing its participants with high-quality, unbiased and state-of-the-art education. We assess potential conflicts of interest with faculty, planners and managers of CME activities. Real or apparent conflicts of interest are identified and resolved through a conflict of interest resolution process. In addition, all activity content is reviewed by both a member of the RTP scientific staff and an external, independent physician reviewer for fair balance, scientific objectivity of studies referenced and patient care recommendations.

**FACULTY** — The following faculty (and their spouses/partners) reported real or apparent conflicts of interest, which have been resolved through a conflict of interest resolution process:

Hagop M Kantarjian, MD  
Chairman and Professor, Leukemia Department  
The University of Texas MD Anderson Cancer Center  
Houston, Texas

No real or apparent conflicts of interest to disclose.

David L Porter, MD  
Abramson Cancer Center  
University of Pennsylvania Health System  
Jodi Fisher Horowitz Professor of Leukemia Care Excellence  
Director, Blood and Marrow Transplantation  
Philadelphia, Pennsylvania

Contracted Research: Novartis Pharmaceuticals Corporation;  
Other Remunerated Activities: Genentech BioOncology (faculty and spouse).

**EDITOR** — Dr Love is president and CEO of Research To Practice, which receives funds in the form of educational grants to develop CME activities from the following commercial interests: AbbVie

Inc, Algeta US, Amgen Inc, Astellas, AstraZeneca Pharmaceuticals LP, Aveo Pharmaceuticals, Bayer HealthCare Pharmaceuticals, Biodesix Inc, Biogen Idec, Boehringer Ingelheim Pharmaceuticals Inc, Bristol-Myers Squibb Company, Celgene Corporation, Daiichi Sankyo Inc, Dendreon Corporation, Eisai Inc, Exelixis Inc, Genentech BioOncology, Genomic Health Inc, Gilead Sciences Inc, Incyte Corporation, Lilly, Medivation Inc, Merck, Millennium: The Takeda Oncology Company, Novartis Pharmaceuticals Corporation, Novocure, Onyx Pharmaceuticals Inc, Prometheus Laboratories Inc, Regeneron Pharmaceuticals, Sanofi, Seattle Genetics, Spectrum Pharmaceuticals Inc, Teva Oncology and VisionGate Inc.

**RESEARCH TO PRACTICE STAFF AND EXTERNAL REVIEWERS** — The scientific staff and reviewers for Research To Practice have no real or apparent conflicts of interest to disclose.

This educational activity contains discussion of published and/or investigational uses of agents that are not indicated by the Food and Drug Administration. Research To Practice does not recommend the use of any agent outside of the labeled indications. Please refer to the official prescribing information for each product for discussion of approved indications,

contraindications and warnings. The opinions expressed are those of the presenters and are not to be construed as those of the publisher or grantors.

This activity is supported by educational grants from Boehringer Ingelheim Pharmaceuticals Inc, Celgene Corporation, Genentech BioOncology/Biogen Idec, Millennium: The Takeda Oncology Company, Onyx Pharmaceuticals Inc, Seattle Genetics and Spectrum Pharmaceuticals Inc.

Hardware/Software Requirements:

A high-speed Internet connection

A monitor set to 1280 x 1024 pixels or more

Internet Explorer 7 or later, Firefox 3.0 or later, Chrome, Safari 3.0 or later

Adobe Flash Player 10.2 plug-in or later

Adobe Acrobat Reader

(Optional) Sound card and speakers for audio

Last review date: May 2014

Expiration date: May 2015

To go directly to slides and commentary for this issue, [click here](#).

On this final issue of our review of select key papers presented at the American Society of Hematology annual meeting, we focus on a handful of fascinating early clinical reports on CART and a smattering of what's new in AML.

### **CART clinical trial data**

Although we heard about the seemingly miraculous effects of this novel therapeutic approach in 2012 in Atlanta, New Orleans was the true coming out party for CART-based approaches, specifically those targeting CD19. Led by the powerhouse team at the University of Pennsylvania, which includes Dr David Porter, we were treated to numerous fascinating presentations that have generated great excitement and enthusiasm.



**David L Porter, MD**

Since ASH I have been fortunate enough to interview Dr Porter on 2 occasions ([click for audio](#)), and it is impossible to hear about this dramatic story without getting goose bumps.

The concept behind Penn's CART-based approach is intriguing. Patients undergo leukapheresis, after which their T-cells are transduced with a lentivirus encoding an anti-CD19 single-chain variable fragment linked to 4-1BB and CD3- $\zeta$  signaling domains. The genetically modified cells are then expanded ex vivo, and soon after a course of lymphocyte-depleting chemotherapy they are then reinfused into the patient where, as documented previously and in new reports at ASH, they expand (up to 10,000-fold), persist functionally (beyond 3 years) and exert a direct antitumor effect.

Early efforts by the group have focused on very advanced chronic lymphocytic leukemia (CLL) and B-cell acute lymphoblastic leukemia (ALL), and although they hoped to see some discernible benefit in early testing, as related by Dr Porter, the initial clinical responses were stunning in rapidity and depth. Much of this work was updated and expanded on in New Orleans. Dr Stephan Grupp [presented longer follow-up outcomes](#) from 17 adults and children with relapsed/refractory ALL revealing that 82% achieved a complete response (CR). Similarly, Dr Porter [provided an update](#) from their Phase I study in which 8 of 14 patients with extensively pretreated CLL had objective responses, including 4 patients with CRs, none of whom have yet experienced

relapse. He also unveiled data from their dose-finding randomized Phase II study demonstrating that 7 of 18 patients responded, including 3 CRs, with no correlation between dose and outcome or toxicity.

Perhaps the highlight of these presentations, at least in my mind, was an impressive set of scans that Dr Porter displayed from a patient with del(17p) CLL treated on their original pilot study. Amazingly, 3 months after receiving the CART infusion this individual, whose disease had progressed through 10 prior therapies, including ibrutinib and radiation therapy, was pretty much disease free in peripheral blood and bone marrow.

In discussing this work with Dr Porter, I was eager to learn more about the profound and rapid tumor lysis/cytokine release syndrome (CRS) that has been described with this therapy. I was wide-eyed as he detailed the team's early experiences with this scary complication that generally occurred within the first few weeks of treatment at the peak of initial T-cell expansion and initially led to life-threatening multiorgan failure. What was perhaps most compelling was that the team was able to quickly determine that the key cytokine causing this syndrome was IL-6, and they were able to successfully intervene with a novel IL-6 receptor antibody approved for rheumatoid arthritis (tocilizumab).

Penn is not alone at the forefront of this research, and at ASH the Memorial group also reported activity with a slightly different anti-CD19 CAR platform across several diseases. In high-risk CLL, Dr Jae Park **presented preliminary data** from a Phase I study evaluating this approach as consolidation after up-front rituximab-chemotherapy. While the data set was quite small — 8 patients with 2 CRs and 2 partial responses — it provides an important proof of principal that ultimately may allow some patients with CLL to receive a short-term treatment that will lead to prolonged tumor control. Relevantly, the study results suggest greater benefit in patients with lower tumor burden, and Dr Porter believes that an important future strategy will be initial cytoreduction, particularly with novel B-cell inhibitors like ibrutinib, followed by an attempt at cure with CART.

Memorial's Dr Marco Davila **also reported** on CAR therapy as a bridge to allotransplant in patients with B-cell ALL, including those with Philadelphia chromosome-positive disease. Importantly, responses were rapid, occurring as early as 7 to 14 days, CRS was manageable and 10 of 12 patients with detectable disease before CAR therapy became minimal residual disease-negative. Four patients went on to allotransplant with 5 more being prepped for it.

The NCI, too, is very involved in this field, and at ASH they provided us with more data on the **use of this technology in ALL** as well as our **first look at it in B-cell lymphomas**, where 5 of 8 patients with chemotherapy-refractory diffuse large B-cell lymphoma or primary mediastinal B-cell lymphoma experienced objective tumor responses.

The next step for this fascinating and innovative treatment strategy is to not only obtain more data but also to investigate the feasibility and reproducibility of doing this on a larger-scale basis.

## AML: Life beyond 3 plus 7

The dismal current landscape of this important cause of mortality is reflected by the fact that the only compound approved for this disease in the last 20 years, gemtuzumab ozogamicin (GO), was removed from the market by the FDA 4 years ago due to toxicity concerns.

As such, for far too long conversations about AML management have revolved around optimal induction and consolidation chemotherapy doses and schedules and the role for various transplant strategies. However, the rapidly emerging translational and related clinical science that permeates ongoing AML research has many optimistic that brighter days are ahead ([click for summary slides of studies discussed below](#)).

One of the more talked about areas is management of the 30% of patients with FMS-like tyrosine kinase 3 (FLT3) mutations, and in the Big Easy Dr Jorge Cortes presented a provocative study on the effects of the as yet unapproved FLT3 inhibitor quizartinib. One of the proposed benefits of this agent is that it is more specific for the target than tyrosine kinase inhibitors such as sorafenib, which is sometimes employed off label when no other alternatives exist. In this Phase II study using lower doses of quizartinib in 76 patients, a 47% CR rate with acceptable toxicities was reported. All eyes are on the ongoing Phase III study evaluating this compound in hopes that it may end up as a useful tool in practice.

The MD Anderson group also reported on SGI-110 — a novel molecule that combines decitabine with guanosine to produce a longer half-life and potentially higher areas under the curve than decitabine. Importantly, in this Phase II study this subcutaneously administered hypomethylating agent resulted in an encouraging preliminary objective response rate of 53% in elderly patients with treatment-naïve AML. Dr Hagop Kantarjian and his group at MD Anderson are interested in doing a study comparing this agent to conventional decitabine or azacitidine.

Because activating KIT mutations are present in 25% to 30% of patients with core binding factor (CBF) AML, it has been hypothesized that KIT inhibition might provide therapeutic benefit. In this regard, the CALGB reported on a single-arm trial evaluating the addition of dasatinib to induction chemotherapy in patients with molecular confirmation of CBF AML. Reported at ASH, the trial resulted in an encouraging CR rate of 92% in 59 evaluable patients, but many are reserving judgment about this approach until further follow-up is available.

Given its activity in a multitude of hematologic cancers, it should probably come as no surprise that lenalidomide is also being evaluated in AML. Significantly, preliminary results look encouraging with a CR rate of 43% in 37 elderly patients over age 70 with low-dose lenalidomide added to low-dose Ara-C (LDAC). Perhaps even more importantly, a 5-gene molecular signature has been identified that appears highly predictive of treatment response with 87% overall accuracy.

Additionally, although the previously mentioned anti-CD33 antibody-drug conjugate GO is no longer available, at ASH we saw data reinforcing its evidence-based benefit most

specifically for patients with CBF AML. Described in [a meta-analysis, a pediatric study and a longitudinal analysis](#) of trials of the UK MRC/NCRI group, these results all point to a modest advantage with limited toxicity. Whether GO will make it back into the clinic, however, remains unclear.

Finally, in terms of the FDA and AML, one of the more interesting and positive recent developments from the agency has been the 2012 implementation of the “Breakthrough Therapy” designation pathway, which fast-tracks promising agents in diseases with important unmet needs. Over the past year on our CME programs we have discussed many exciting oncology compounds that have earned this designation, and just 2 weeks ago we saw this novel pathway in action as the second-generation ALK inhibitor ceritinib was granted accelerated approval in ALK-positive non-small cell lung cancer based on the results of a single-arm, open-label clinical trial enrolling 163 patients.

Last fall, the selective and potent polo-like kinase (PLK) inhibitor volasertib became the only “AML drug” to join this short “Breakthrough” list. Volasertib inhibits PLK1 — the best characterized of the 5 known human PLKs and a critical enzyme regulating mitosis — resulting in cell cycle arrest and ultimately cell death (apoptosis). At ASH 2012 a randomized Phase I/II trial reported an impressive response rate advantage when the agent was added to LDAC in patients not eligible for intensive induction. These encouraging results led to the ongoing Phase III POLO-AML-2 trial with a similar randomization in patients age 65 or older with previously untreated AML not eligible for intensive induction. Although the future of this interesting agent is unclear, it seems plausible that in the next few years AML will join the other myeloid cancers in seeing the documented benefit of important new and clinically useful treatment strategies based on the evolution of understanding the underlying disease biology.

This concludes our ASH review series. If you’re heading to Chicago this month, join us at the end of each day for a series of evening symposia as we review what’s happening in [lung cancer](#), [gastrointestinal cancers](#), [non-Hodgkin lymphoma/multiple myeloma](#) and [HER2-positive breast cancer](#).

Neil Love, MD

[Research To Practice](#)

Miami, Florida

Research To Practice is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Research To Practice designates this enduring material for a maximum of 2 *AMA PRA Category 1 Credits™*. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

This activity is supported by educational grants from Boehringer Ingelheim Pharmaceuticals Inc, Celgene Corporation, Genentech BioOncology/Biogen Idec, Millennium: The Takeda Oncology Company, Onyx Pharmaceuticals Inc, Seattle Genetics and Spectrum Pharmaceuticals Inc.



# Reinduction of Complete Remissions Using 19-28z CAR T-Cell Therapy in Relapsed B-Cell ALL

## Presentation discussed in this issue

Davila M et al. **Safe and effective re-induction of complete remissions in adults with relapsed B-ALL using 19-28z CAR CD19-targeted T cell therapy.** *Proc ASH 2013*; **Abstract 69**.

Slides from a presentation at ASH 2013 and transcribed comments from recent interviews with Hagop M Kantarjian, MD (1/29/14) and David L Porter, MD (3/3/14)

## Safe and Effective Re-Induction of Complete Remissions in Adults with Relapsed B-ALL Using 19-28z CAR CD19-Targeted T Cell Therapy

**Davila ML et al.**

*Proc ASH 2013*; Abstract 69.

Research  
To Practice®



## Background

- A majority of adults with B-cell acute lymphoblastic leukemia (B-ALL) relapse and develop chemorefractory disease.
- Novel, nonchemotherapy-based treatments are needed for this group of patients.
- Minimal residual disease (MRD)-negative remissions were observed in small numbers of patients with B-ALL treated with adoptive immunotherapy using T cells genetically engineered to express chimeric antigen receptors (CARs) targeting CD19, a molecule expressed on normal and malignant B cells (*Sci Transl Med* 2013;5(177):177ra38).
- **Study objective:** To determine the safety, appropriate dose and efficacy of CARs in adults with relapsed/refractory (RR) B-ALL.

Davila ML et al. *Proc ASH* 2013;Abstract 69 (abstract only).

Research  
To Practice®

## Phase I Study Methods (NCT01044069)

- Adults with RR CD19+ B-ALL (target enrollment = 40) were enrolled.
- Patients were leukapheresed and reinduced with salvage chemotherapy.
- Patients received an infusion of the genetically modified T cells ( $3 \times 10^6$  19-28z CAR T cells/kg) after conditioning chemotherapy.
  - 19-28z is composed of a CD19 binding domain fused to the signaling domains of the CD28 costimulatory receptor and the  $\zeta$  chain of the CD3 complex.
- 13 patients enrolled to date: 11 had RR disease, 2 were enrolled and leukapheresed during CR1 but were not treated until relapse.

Davila ML et al. *Proc ASH* 2013;Abstract 69 (abstract only).

Research  
To Practice®

## Baseline Characteristics

- Median patient age was 42 (range: 23 to 74).
- 3 of the 13 patients had Philadelphia chromosome-positive (Ph+) B-ALL.
- The required T-cell dose was achieved in all but 1 patient.
- 7 of the 13 patients were infused with 19-28z CAR T cells while they had gross residual disease (>5% to 70% blasts in the bone marrow).
- The remaining patients had MRD, detected by flow cytometry or deep sequencing, at the time of 19-28z CAR T-cell infusion.

Davila ML et al. *Proc ASH 2013*;Abstract 69 (abstract only).

Research  
To Practice®

## Clinical Outcomes

- 10 of the 12 patients with detectable disease before T-cell infusion developed MRD-negative responses.
  - 5 patients with gross residual disease (blasts >5% in bone marrow) became MRD-negative.
  - 5 MRD-positive patients became MRD-negative.
- The rapidity of the responses was remarkable, with MRD-negative results obtained as early as 7 to 14 days after T-cell infusion.
- Patients with relapsed Ph+ B-ALL achieved MRD-negative status as determined by both deep sequencing for the IgH rearrangement and qPCR for the BCR-ABL transcript.

Davila ML et al. *Proc ASH 2013*;Abstract 69 (abstract only).

Research  
To Practice®



## Adverse Events

- 6 of 13 patients developed toxicities, including high-grade fevers, hypotension, hypoxia, mental status changes and seizures.
- These episodes occurred for approximately 1 week before they were halted by treatment with steroids or tocilizumab.
- All patients completely recovered and were able to leave the hospital.
- The occurrence of toxicities correlated with tumor burden such that patients with gross residual disease (>5% blasts in bone marrow) developed toxicities, whereas those with MRD had no evidence of toxicities.
- No toxicities were experienced by the other 7 patients.

Davila ML et al. *Proc ASH 2013*;Abstract 69 (abstract only).

Research  
To Practice®

## Author Conclusions

- The potent induction of MRD-negative responses and the reversibility of toxicities occurring in a subset of patients strongly support integration of this therapy into the modern paradigm for B-ALL therapy.
- Of the 13 patients treated, 4 underwent an allogeneic stem cell transplant (allo-SCT) and 5 are being prepared for an allo-SCT. Of the remaining 4 patients, 1 is a nonresponder, 1 relapsed, 1 is in CR2 (medically not eligible for allo-SCT) and 1 has not been evaluated for response yet.
- Toxicities associated with this therapy are predictable and manageable.
- The remarkable MRD-negative reinduction rate and facilitation for allo-SCT reported warrant further evaluation of this therapy in a Phase II protocol.

Davila ML et al. *Proc ASH 2013*;Abstract 69 (abstract only).

Research  
To Practice®

### **Investigator Commentary: Reinduction of Complete Remissions in Relapsed B-ALL Using 19-28z CAR T-Cell Therapy**

In this study 10 of the 12 patients with relapsed/refractory B-ALL treated with CAR therapy achieved MRD-negative status. In terms of toxicity, most patients develop delayed cytokine release syndrome after a week or 2. It manifests as fever, bone and muscle aches, nausea and vomiting and anorexia. These side effects can be effectively managed with steroids and anti-IL-6.

This is an exciting approach that needs further study. I believe that it is possible to develop this therapy for use in the front-line setting. Patients with ALL who have MRD could be treated with CAR therapy, potentially increasing the chance of a cure and obviating the need for additional chemotherapy.

***Interview with Hagop M Kantarjian, MD, January 29, 2014***

Research  
To Practice®

### **Investigator Commentary: Reinduction of Complete Remissions in Relapsed B-ALL Using 19-28z CAR T-Cell Therapy (Continued)**

This study used a CAR construct that is slightly different than the one used in our studies, but it appears to have extremely potent activity in ALL. The authors reported that 10 of 12 patients with detectable disease prior to T-cell infusion subsequently achieved MRD-negative status. The highlight of the study was that the therapy was effective even in patients with Ph+ B-ALL, which is important and unique. Four of the 13 patients who underwent treatment were able to undergo allo-SCT and another 5 patients were being prepared for transplant. The investigators were using this approach as a bridge to transplant for patients who would have been otherwise ineligible so that they might have a chance of being cured.

***Interview with David L Porter, MD, March 3, 2014***

Research  
To Practice®